


HIV Surveillance in Women

National Center for HIV/AIDS, Viral Hepatitis, STD & TB Prevention
Division of HIV/AIDS Prevention



Diagnoses of HIV Infection among Adult and Adolescent Females, by Race/Ethnicity, 2009–2013 United States and 6 Dependent Areas



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection among Adult and Adolescent Females, by Race/Ethnicity, 2013—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	49	5.1
Asian	159	2.2
Black/African American	5,867	34.8
Hispanic/Latino ^a	1,419	7.0
Native Hawaiian/other Pacific Islander	15	7.3
White	1,579	1.8
Multiple races	190	9.3
Total^b	9,278	6.9

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^a Hispanics/Latinos can be of any race.


^b Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



Diagnoses of HIV Infection and Population among Adult and Adolescent Females, by Race/Ethnicity 2013—United States

Diagnoses of HIV Infection


N=9,278



American Indian/Alaska Native
Asian
Black/African American

Female Population

N = 134,687,181



Hispanic/Latino^a
Native Hawaiian/other Pacific Islander
White


Multiple races

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.


^a Hispanics/Latinos can be of any race.

Diagnoses of HIV Infection among Adult and Adolescent Females, by Race/Ethnicity and Transmission Category 2013—United States and 6 Dependent Areas



Black/African American
N=5,876



Hispanic/Latino^a
N=1,610



White
N=1,579



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.

^b Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^c Includes blood transfusion, perinatal exposure, and risk factor not reported or not identified.

Diagnoses of HIV Infection among Adult and Adolescent Females, by Transmission Category and Age at Diagnosis 2013—United States and 6 Dependent Areas

Transmission category	Age at Diagnosis (in years)				
	13–19	20–24	25–34	35–44	≥45
	N=376	N=923	N=2,378	N=2,257	N=3,544
	%	%	%	%	%
Injection drug use	6.1	9.8	11.6	12.2	14.8
Heterosexual contact ^a	84.0	87.7	87.6	87.6	84.9
Other ^b	9.9	2.5	0.8	0.2	0.3
Total	100.0	100.0	100.0	100.0	100.0

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.


^b Includes blood transfusion, perinatal exposure, and risk factor not reported or not identified.



Rates of Diagnoses of HIV Infection among Adult and Adolescent Females, 2013—United States and 6 Dependent Areas


N = 9,479

Total rate = 6.9



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

Diagnoses of HIV Infection among Adult and Adolescent Females, by Region and Race/Ethnicity 2013—United States



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.

Deaths of Adult and Adolescent Females with Diagnosed HIV Infection, by Race/Ethnicity 2012—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	22	2.3
Asian ^a	4	0.1
Black/African American	2,782	16.7
Hispanic/Latino ^b	625	3.2
Native Hawaiian/other Pacific Islander	0	0.0
White	866	1.0
Multiple races	222	11.2
Total^c	4,521	3.4

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Deaths of persons with diagnosed HIV infection may be due to any cause. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.


^c Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



Rates of Adult and Adolescent Females Living with Diagnosed HIV Infection, Year-end 2012—United States and 6 Dependent Areas


N = 229,176

Total rate = 169.4




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

Stage 3 (AIDS) Classifications among Adult and Adolescent Females, 1985–2013—United States and 6 Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

Stage 3 (AIDS) Classifications among Adult and Adolescent Females, by Race/Ethnicity and Year of Diagnosis 1985–2013—United States and 6 Dependent Areas




Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.

^b Includes Asian/Pacific Islander legacy cases.

Percentages of Stage 3 (AIDS) Classifications among Adult and Adolescent Females, by Race/Ethnicity and Year of Diagnosis 1985–2013—United States and 6 Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.

^b Includes Asian/Pacific Islander legacy cases.



Stage 3 (AIDS) Classifications among Adult and Adolescent Females, by Race/Ethnicity, 2013—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	25	2.6
Asian ^a	60	0.8
Black/African American	4,238	25.1
Hispanic/Latino ^b	891	4.4
Native Hawaiian/other Pacific Islander	7	3.5
White	971	1.1
Multiple races	231	11.3
Total^c	6,424	4.8

Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.

^c Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



Stage 3 (AIDS) Classifications among Adult and Adolescent Females, by Transmission Category and Age at Diagnosis 2013—United States and 6 Dependent Areas

Age at Diagnosis (in years)

	13–19	20–24	25–34	35–44	≥45
Transmission category	N=147	N=348	N=1,308	N=1,821	N=2,915
	%	%	%	%	%
Injection drug use	2.4	5.2	13.0	16.4	23.1
Heterosexual contact ^a	39.3	79.0	85.4	83.4	76.6
Other ^b	58.3	15.8	1.6	0.2	0.3
Total	100.0	100.0	100.0	100.0	100.0

Note. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.


^b Includes blood transfusion, perinatal exposure, and risk factor not reported or not identified.



Rates of Stage 3 (AIDS) Classifications among Adult and Adolescent Females with HIV Infection, 2013—United States and 6 Dependent Areas


N=6,537

Total Rate = 4.8



Note: All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

Stage 3 (AIDS) Classifications among Adult and Adolescent Females, by Region and Race/Ethnicity 2013—United States




Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.



Rates of Adult and Adolescent Females Living with Diagnosed HIV Infection Ever Classified as Stage 3 (AIDS) Year-end 2012—United States and 6 Dependent Areas

N=123,320 Total rate = 91.1



Note: All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.